arXiv:0706.3782v2 [astro-ph] 5 Jul 2007

The Virgo - MiniGRAIL cross correlation for the detection of scalar gravitational waves

Christian Corda

February 1, 2008

INFN - Sezione di Pisa and Università di Pisa, Via F. Buonarroti 2, I - 56127
PISA, Italy

E-mail address: christian.corda@ego-gw.it

Abstract

After a review of the frequency - dependent angular pattern of interferometers in the TT gauge for scalar gravitational waves (SGWs), which has been recently analysed by Capozziello and Corda, in this letter the result is used to study the cross correlation between the Virgo interferometer and the MiniGRAIL resonant sphere. It is shown that the overlap reduction function for the cross correlation bewteen Virgo and the monopole mode of MiniGRAIL is very small, but a maximum is also found in the correlation at about $2710Hz$, in the range of the MiniGRAIL sensitivity.

PACS numbers: 04.80.Nn, 04.30.Nk, 04.50.+h

The design and construction of a number of sensitive detectors for gravitational waves (GWs) is underway today. There are some laser interferometers and many bar detectors currently in operation and several interferometers and bars are in a phase of planning and proposal stages (for the current status of gravitational waves experiments see [1, 2, 3, 4, 5]).

The results of these detectors will have a fundamental impact on astrophysics and gravitation physics. There will be lots of experimental data to be analyzed, and theorists will be forced to interact with lots of experiments and data analysts to extract the physics from the data stream.

Detectors for GWs will also be important to verify that GWs only change distances perpendicular to their direction of propagation and to confirm or ruling out the physical consistency of General Relativity or of any other theory of gravitation [6, 7, 8, 9, 10].

These detectors are in principle sensitive also to a hypotetical *scalar* component of gravitational radiation, that appears in extended theories of gravity like scalar-tensor gravity [11, 12, 13, 14] and high order theories [7, 8, 9, 10].

While the response of interferometers to tensorial waves has been calculated in lots of works (see for example [15, 16, 17]), the coupling between interferometers and scalar waves is a more recent field of interest and it has not an analogous number of works in the literature. Only recently, Capozziello and Corda have shown that the response of an interferometer to massless SGWs is invariant in three different gauges known in literature in its full frequency dependence [6] while previous works started from the assumption that the wavelength of the SGW is much larger than the distance between the test masses (see [12, 13, 14] for examples).

In this letter, after a review of the frequency - dependent angular pattern of interferometers in the TT gauge, which is due to provide a context to bring out the relevance of the argument, the result will be used to study a cross correlation between the Virgo interferometer and the MiniGRAIL resonant sphere [18, 19] generalizing the analysis of [13] for a real detector.

Following [6], in the TT gauge, for a purely massless SGW propagating in the positive z direction, with the interferometer located at the origin of the coordinate sistems with arms in the $\overrightarrow{x}$ and $\overrightarrow{y}$ directions, the metric perturbation is given by (see [6, 12, 13], and note that we work with $c = 1$ and $\hbar = 1$ in this letter):

$$h_{\mu\nu}(t-z) = \Phi(t-z)e_{\mu\nu}^{(s)}, \tag{1}$$

where $\Phi \ll 1$, $e_{\mu\nu}^{(s)} \equiv diag(0,1,1,0)$, and the line element is

$$ds^2 = -dt^2 + dz^2 + [1 + \Phi(t-z)][dx^2 + dy^2]. \tag{2}$$

To compute the response function for an arbitrary propagating direction of the SGW one recalls that the arms of the interferometer are in the $\overrightarrow{u}$ and $\overrightarrow{v}$ directions, while the x, y, z frame of (2) is adapted to the propagating SGW. Thus a spatial rotation of the coordinate system is needed:

$$\begin{array}{ccc} u & = & -x\cos\theta\cos\phi + y\sin\phi + z\sin\theta\cos\phi \\ \\ v & = & -x\cos\theta\sin\phi - y\cos\phi + z\sin\theta\sin\phi \\ \\ w & = & x\sin\theta + z\cos\theta, \end{array} \tag{3}$$

or, in terms of the x, y, z frame:



Figure 1: a SGW propagating from an arbitrary direction

$$\begin{aligned} x &= -u\cos\theta\cos\phi - v\cos\theta\sin\phi + w\sin\theta \\ y &= u\sin\phi - v\cos\phi \\ z &= u\sin\theta\cos\phi + v\sin\theta\sin\phi + w\cos\theta. \end{aligned} \tag{4}$$

In this way the SGW is propagating from an arbitrary direction to the interferometer (see figure 1).

The beam splitter is also put in the origin of the new coordinate system (i.e. $u_b = 0$, $v_b = 0$).

The transformation for the metric tensor is [20]:

$$g^{ik} = \frac{\partial x^i}{\partial x'^l}\frac{\partial x^k}{\partial x'^m} g'^{lm}. \tag{5}$$

By using eq. (3), eq. (4) and eq. (5), the line element (2) in the $\overrightarrow{u}$ direction becomes:

$$ds^2 = -dt^2 + [1 + (1 - \sin^2\theta\cos^2\phi)\Phi(t - u\sin\theta\cos\phi)]du^2. \tag{6}$$

A good way to analyze variations in the proper distance (time) is by means of “bouncing photons” (see [6, 17] and figure 2).



Figure 2: photons can be launched from the beam-splitter to be bounced back by the mirror

Let us compute the variaton of the proper distance that a photon covers to make a round-trip from the beam-splitter to the mirror of an interferometer with the line-element choice (6).

A special property of the TT gauge is that an inertial test mass initially at rest in these coordinates, remains at rest throughout the entire passage of the SGW [6, 12, 13]. Here we have to clarify the use of words " at rest": we want to mean that the coordinates of the test masses do not change in the presence of the SGW. The proper distance between the beam-splitter and the mirror of the interferometer changes even though their coordinates remain the same [6, 13].

Putting the condition for null geodesics $ds^2 = 0$ in eq. (6) the coordinate velocity of the photon, which is a convenient quantity for calculations of the photon propagation time between the beam-splitter and the mirror [6, 17], is:

$$v^2 \equiv (\frac{du}{dt})^2 = \frac{1}{[1+(1-\sin^2\theta\cos^2\phi)\Phi(t-u\sin\theta\cos\phi)]}. \tag{7}$$

Because the coordinates of the beam-splitter $u_b = 0$ and of the mirror $u_m = L$ do not change under the influence of the SGW, the duration of the forward trip can be found as

$$T_1(t) = \int_0^L \frac{du}{v(t'-u\sin\theta\cos\phi)}, \tag{8}$$

with

$$t' = t-(L-u).$$

In the last equation t' is the retardation time (i.e. t is the time at which the photon arrives in the position L, so $L-u = t-t'$).

To first order in Φ this integral is approximated by

$$T_1(t) = T + \frac{1-\sin^2\theta\cos^2\phi}{2}\int_0^L \Phi(t'-u\sin\theta\cos\phi)du, \tag{9}$$

where

$$T = L$$

is the transit time of the photon in the absence of the SGW. Similiary, the duration of the return trip will be

$$T_2(t) = T + \frac{1-\sin^2\theta\cos^2\phi}{2}\int_L \Phi(t'-u\sin\theta\cos\phi)(-du), \tag{10}$$

though now the retardation time is

$$t' = t - u.$$

The round-trip time will be the sum of $T_2(t)$ and $T_1[t-T_2(t)]$. The latter can be approximated by $T_1(t-T)$ because the difference between the exact and the approximate values is second order in Φ. Then, to first order in Φ, the duration of the round-trip will be

$$T_{r.t.}(t) = T_1(t-T) + T_2(t). \tag{11}$$

By using eqs. (9) and (10) we see immediatly that deviations of this round-trip time (i.e. proper distance) from its imperurbated value are given by

$$\begin{gathered}\delta T(t) = \tfrac{1-\sin^2\theta\cos^2\phi}{2}\int_0^L[\Phi(t-2T+u(1-\sin\theta\cos\phi))+ \\ +\Phi(t-u(1+\sin\theta\cos\phi))]du.\end{gathered} \tag{12}$$

Now, using the Fourier transform of the scalar field, defined by

$$\tilde{\Phi}(\omega) = \int_{-\infty}^{\infty} dt\Phi(t)\exp(i\omega t), \tag{13}$$

the analysis can be transled in the frequency domain obtaining

$$\frac{\delta\tilde{T}(\omega)}{T} = \Upsilon_u(\omega)\tilde{\Phi}(\omega) \tag{14}$$

where the total response function in the $\overrightarrow{u}$ direction is given by:

$$\begin{gathered}\Upsilon_u(\omega) = \tfrac{1}{2i\omega L}[-1+\exp(2i\omega L)+ \\ +\sin\theta\cos\phi((1+\exp(2i\omega L)-2\exp i\omega L(1+\sin\theta\cos\phi))].\end{gathered} \tag{15}$$

In the same way the line element (2) in the $\overrightarrow{v}$ direction becomes:

$$ds^2 = -dt^2 + [1+(1-\sin^2\theta\sin^2\phi)\Phi(t-v\sin\theta\sin\phi)]dv^2, \tag{16}$$

thus, with the same analysis of the $\overrightarrow{u}$ direction, the total response function in the $\overrightarrow{v}$ direction is obtained:

$$\begin{array}{c}\Upsilon_v(\omega)=\frac{1}{2i\omega L}[-1+\exp(2i\omega L)+\\+\sin\theta\sin\phi((1+\exp(2i\omega L)-2\exp i\omega L(1+\sin\theta\sin\phi))].\end{array}\tag{17}$$

Then, the total response function of an interferometer, defined by $\tilde{H}(\omega)\equiv\Upsilon_u(\omega)-\Upsilon_v(\omega)$, is:

$$\begin{array}{c}\tilde{H}(\omega)=\frac{\sin\theta}{2i\omega L}\{\cos\phi[1+\exp(2i\omega L)-2\exp i\omega L(1+\sin\theta\cos\phi)]+\\-\sin\phi[1+\exp(2i\omega L)-2\exp i\omega L(1+\sin\theta\sin\phi)]\}.\end{array}\tag{18}$$

Eq. (18) is also in perfect agreement with the detector pattern of [13] and [14] in the low frequencies limit ($\omega\to 0$):

$$\tilde{H}(\omega\to 0)=-\sin^2\theta\cos 2\phi.\tag{19}$$

In figs. 3 and 4 the absolute value of the total response function of the Virgo interferometer ($L=3$ Km)) for SGWs with $\theta=\frac{\pi}{4}$ and $\phi=\frac{\pi}{3}$, and the angular dependence of the response of the Virgo interferometer for a SGW with a frequency of $f=100Hz$ are shown respectively.

Now, the analysis of [13] , where a cross correlation between the Virgo interferometer and a hypothetical resonant sphere in Frascati, near Rome, was studied, will be generalized for a real detector, the MiniGRAIL sphere [18, 19].

In [13] the overlap reduction function for SGWs has been computed generalizing the well known Flanagan's overlap reduction function for ordinary (tensorial) GWs defined in [21]. The overlap reduction function for SGWs has also been computed in [22].

In the cross correlation between a sphere and an interferometer, the monopole mode of a resonant sphere is especially interesting, because it cannot be excited by tensorial waves [13, 22].

The authors of [13] found the relation (see eq. (56) of [13])

$$\Gamma(f)=(\sin^2\theta\cos 2\phi)j_2(2\pi fd),\tag{20}$$

where j_2 is the second spherical Bessel function and θ and ϕ are the angular coordinates of the MINIGRAIL sphere with respect the Virgo interferometer. (see figure 5).

In eq. (20) the low frequencies approximation (19) has been used. Replacing the low frequencies approximation with our frequency dependent angular pattern (18) the more general overlap reduction function is obtained:

$$\begin{array}{c}\Gamma(f)=\frac{-\sin\theta}{4i\pi fL}\{\cos\phi[1+\exp(4i\pi fL)-2\exp i2\pi fL(\sin\theta\cos\phi-1)]+\\-\sin\phi[1+\exp(4i\pi fL)-2\exp i2\pi fL(\sin\theta\sin\phi-1)]\}j_2(2\pi fd).\end{array}\tag{21}$$



Figure 3: the absolute value of the total response function of the Virgo interferometer to SGWs for $\theta = \frac{\pi}{4}$ and $\phi = \frac{\pi}{3}$.



Figure 4: the angular dependence of the response of the Virgo interferometer for a SGW with a frequency of $f = 100Hz$



Figure 5: The Virgo-MiniGRAIL cross correlation



Figure 6: the absolute value of the total response function of the Virgo interferometer to SGWs for the Virgo- MiniGRAIL direction.

MiniGRAIL works in the frequency range between $2670-3130Hz$ [18, 19] with a resonance frequency of about $2900Hz$, and, because the angular pattern in the Virgo- MiniGRAIL direction is almost constant in this range (figure 6), the function $j_2(2\pi fd)$ is a good approximation for the frequency dependence of the overlap reduction function.

In figure 7 the frequency dependence of the overlap reduction function for the approximate distance between the location of the Virgo interferometer and the location of the MiniGRAIL resonant sphere in the MiniGRAIL frequency range is shown. The location of MiniGRAIL is Lat. N 52.16, Lon E 4.48, while the location of Virgo is Lat. N 43.63, Lon E 10.50 and the Virgo-MiniGRAIL distance is about 1090 kms. Figure 7 shows that the overlap reduction function for the Virgo-MiniGRAI cross correlation is very small, but there is a maximum at about $2710Hz$. Approximately this maximum is reached when

$$(\frac{f}{2710Hz})(\frac{d}{1090km})\simeq 1. \tag{22}$$

Reasuming, in this letter the frequency - dependent angular pattern of in-



Figure 7: the frequency dependence of the Virgo-MiniGRAIL overlap reduction function in the MiniGRAIL frequency range

terferometers in the TT gauge has been reanalyzed following [6]. After this, the result has been used to study a cross correlation between the Virgo interferometer and the MiniGRAIL resonant sphere generalizing the analysis of [13] for a real detector. It has been shown that the overlap reduction function for the cross correlation bewteen Virgo and the monopole mode of MiniGRAIL is very small, but a maximum in the correlation has also be found at about $2710Hz$, in the range of the MiniGRAIL sensitivity.

Acknowledgements

I would like to thank Salvatore Capozziello, Giancarlo Cella and Francesco Rubanu for helpful advices during my work. I have to thank the European Gravitational Observatory (EGO) consortium for the using of computing facilities.

References

[1] Acernese F et al. (the Virgo Collaboration) - Class. Quant. Grav. **23** 19 S635-S642 (2006);

[2] Corda C - Astropart. Phys. **27,** No 6, 539-549 (2007);

[3] Hild S (for the LIGO Scientific Collaboration) - Class. Quant. Grav. **23** 19 S643-S651 (2006)

[4] Willke B et al. - Class. Quant. Grav. **23** 8S207-S214 (2006);

[5] Tatsumi D, Tsunesada Y and the TAMA Collaboration - Class. Quant. Grav. **21** 5 S451-S456 (2004)

[6] Capozziello S and Corda C - Int. J. Mod. Phys. D **15** 1119 -1150 (2006)

[7] Capozziello S - *Newtonian Limit of Extended Theories of Gravity* in *Quantum Gravity Research Trends* Ed. A. Reimer, pp. 227-276 Nova Science Publishers Inc., NY (2005) arXiv:gr-qc/0412088

[8] Capozziello S and Troisi A Phys. Rev. D **72** 044022 (2005)

[9] Allemandi G, Capone M, Capozziello S and Francaviglia M - Gen. Rev. Grav. **38** 1 (2006)

[10] Allemandi G, Francaviglia M, Ruggiero ML and Tartaglia A - Gen. Rev. Grav. **37** 11 (2005)

[11] Brunetti M, Coccia E, Favone V and Fucito F - Phys. Rew. D **59** 044027 (1999)

[12] Tobar ME, Suzuki T and Kuroda K Phys. Rev. D 59 102002 (1999);

[13] Maggiore M and Nicolis A - Phys. Rev. D **62** 024004 (2000); also in gr-qc/9907055

[14] Bonasia N and Gasperini G Phys. Rew. D **71** 104020 (2005)

[15] Misner CW, Thorne KS and Wheeler JA - “Gravitation” - W.H.Feeman and Company - 1973

[16] Maggiore M- Physics Reports **331**, 283-367 (2000)

[17] Rakhmanov M - Phys. Rev. D **71** 084003 (2005)

[18] De Waard A et al - Proceedings of the 6th Edoardo Amalfi Conference ooon Gravitatinal Waves, Bankolu Shinryoukan Kise Nago, Okinawa Japan, June 20-24, 2005

[19] De Waard A et al - Classical and Quantum Gravity **22** S215-S219

[20] Landau L and Lifsits E - “Teoria dei campi” - Editori riuniti edition III (1999)

[21] Flanagan E - Phys. Rev. D **48** 2389 (1993)

[22] Babusci D, Baiotti L, Fucito F and Nagar A - Phys. Rew. D **64** 062001 (2001)